

SYNTHESIS, CHARACTERIZATION OF A NEW COMPOUND 22-(3,5-BIS(3,5,5,6-TETRAMETHYL-1,1-DIOXIDO-5,6-DIHYDRO-2H-1,2-THIAZIN-2-YL)PHENYL)-3,5,5,6-TETRAMETHYL-5,6-DIHYDRO-2H-1,2-THIAZINE 1,1-DIOXIDE, AND STUDY OF ITS BACTERIAL AND FUNGAL ACTIVITY AND MOLECULAR DOCKING

Marwan M. Farhan

University Of Anbar, College of Applied Sciences, Hit, Iraq
mw_mw_888@uoanbar.edu.iq

Received 17.01.2025

Accepted 11.02.2025

Abstract: A synthesis of a new compound (II) with three groups of 1,2-thiazine 1,1-dioxide, called (2-(3,5-bis(3,5,5,6-tetramethyl-1,1-dioxido-5,6-dihydro-2H-1,2-thiazin-2-yl)phenyl)-3,5,5,6-tetramethyl-5,6-dihydro-2H-1,2-thiazine 1,1-dioxide), was synthesized by reacting moles (1:3) of each of benzene-1,3,5-triamine with 3,4,4,6-tetramethyl-3,4-dihydro-1,2-oxathiine 2,2-dioxide (I), respectively. The last compound (I) was prepared by equal moles of each of (4-methylpent-3-en-2-one) with (sulfonylthane). The structural formula of the resulting compounds was confirmed by their spectroscopic identification (FT-IR, ¹H-NMR, ¹³C-NMR). The effect of these two compounds at different concentrations (0.01, 0.1, 1.0, and 10 mg/ml) on the growth of six types of Gram-negative and positive bacteria (*Protans vulgarisginosa*, *Staphylococcus aureus*, *Pseudonas aeruginosa*, *Bacillus subtilus*, and *Enterbacter sp.*) was studied. Compared with the pharmaceutical compound Chloramphenicol, it shows high inhibition values at concentrations (1.0 and 10 mg/ml) and exceeds the reference's inhibition limit. It is also noted that the concentrations of the acid compound (0.01 and 0.1 mg/ml) are acceptable. The effect of these two compounds was also measured using concentrations (10, 25, and 30 mg/ml) on the growth of fungi *A. fumigates*, *T.mentagrophytes*, *C.albicans* and *M. canis zone* (killing). For the concentrations of the prepared compounds, the killing range of compound (II) is much higher compared to compound (I). Furthermore, molecular docking studies have revealed many correlations with amino acid residues in active sites, so these results have the potential to stimulate further research on heterocyclic compounds.

Keywords: Synthesis, new compound, -1,2-thiazine 1,1-dioxide, Dugong, fungi, bacteria

DOI: 10.65382/2221-8688-2026-1-31-41

Introduction

Old, the name 1,4-sultones, 1,4-butanestultone, or δ -sultones was given to the compounds 1,2-oxathiine 2,2-dioxides [1], which are relatively rare compounds and are heterocyclic compounds of which there are multiple isomers at the positions (1,2,3) [2-4]. The least discovered of which is 1,2-oxathiine. A review of the literature shows, that compounds with hexagonal rings containing one or two heteroatoms were prepared (Fig. 1): tetrahydro-1 (1,2-oxathiane 2,2-dioxide), three dihydro-2, 3, 4 isomers, and the completely unsaturated 1,2-oxathiine 2,2-dioxide systems that can be considered as SO₂ analogues of the pyran-2-one system that has been studied in detail and is distinguished by its great biological effect [5-7].

A review of the scientific literature on the

reactions, synthesis, structure, and applications of the 1,2-oxathiine 2,2-dioxide ring system [8] suggests that this heterogeneous ring, which has rarely been studied, can be studied, for example, following traditional and modern synthetic chemistry, as this ring provides great potential as it is considered a basic building block for building many heterogeneous and non-cyclic compounds. Furthermore, δ -sultone is an invaluable reagent, as it is easily soluble in water, as well as in a variety of materials including polymers and dyes [9-13].

These compounds, as well as those containing nitrogen atoms, are also heterogeneous, play an important role in the effect of biological activity, and the use of these compounds appears to be of great importance in

many applications in the field of medicinal chemistry, pharmaceuticals, and agricultural pesticides [14-15].




Fig. 1. Structure and nomenclature of 1,2-oxathiine derivatives

They are used in various compounds with biological properties, including antibacterial [16], anticonvulsant [17], herbicide [18], activity against Gram-positive bacteria [19, 20], anti-HIV [21], anti-inflammatory effects [22], anti-allergic [23], antifungal, and antiviral [24, 25].

They also have pharmaceutical applications due to their stability towards moisture, light, oxygen, and metabolism in the body [26]. They also have wide applications in industry [27], can be used as photosensitizing materials [28], dyes, and agricultural chemicals

[29]. They have a benefit as synthetic aids [30-31] and are used as anti-corrosion agents commercially and in drug discovery [32], materials science [26], and synthetic chemistry [33-35].

The purpose of this study was the synthesis and characterization of a new compound, 22-(3,5-bis(3,5,5,6-tetramethyl-1,1-dioxido-5,6-dihydro-2H-1,2-thiazin-2-yl)phenyl)-3,5,5,6-tetramethyl-5,6-dihydro-2H-1,2-thiazine 1,1-dioxide, and the study of its bacterial and fungal activity and molecular docking.

Experimental part

Synthesis of 3,4,4,6-tetramethyl-3,4-dihydro-1,2-oxathiine 2,2-dioxide (I) [36-38].

The reaction was carried out using the method previously described by Brain [38], with some modifications. 4-Methylpent-3-en-2-one (3.93 g, 0.04 mol) is added gradually to acetic anhydride (45 g), and the mixture is cooled to 0 degrees Celsius. Then, sulfonyl ethane acid (3.68 g, 0.04 mol) is added drop by drop while the reaction temperature remains at 0 degrees Celsius, using an ice bath with salt. The reaction is stirred for two hours, then placed in the refrigerator for 24 hours. Then we separate the reaction mixture by pouring it into an equal amount of water and ice with vigorous stirring. The resulting light yellow precipitate is then separated. A solvent consisting of 80% water and ethanol was used to purify the precipitate by recrystallization. The reaction was followed using the TLC technique. It gave a product (3,4,4,6-tetramethyl-3,4-dihydro-1,2-oxathiine 2,2-dioxide (I)), which has a prismatic shape and is colorless, and its melting point is m.p. 77-

79°C, and yield is 78%. The identity of the resulting compound was confirmed by spectroscopic methods.

Synthesis of 2-(3,5-bis(3,5,5,6-tetramethyl-1,1-dioxido-5,6-dihydro-2H-1,2-thiazin-2-yl)phenyl)-3,5,5,6-tetramethyl-5,6-dihydro-2H-1,2-thiazine 1,1-dioxide (II) [36-38]

Benzene-1,3,5-triamine (0.5 g, 0.004 mol) is gradually added to the compound 3,4,4,6-tetramethyl-3,4-dihydro-1,2-oxathiine 2,2-dioxide (I) (2.28 g, 0.012 mol). In a circular flask equipped with a reflux condenser at a temperature not exceeding 120°C for a period ranging from 40 to 48 hours. After the reaction is over, the mixture is added to dilute hydrochloric acid (1%), then the precipitate is filtered, washed with water, dried, and recrystallized from absolute methanol to give bright white crystal blades with m.p. 122-124 °C and yields of 84%. The identity of the resulting compound was confirmed by spectroscopic methods.

Results and discussion

The derivative of the oxathiine compound was prepared by reacting an equal number of moles of (4-methylpent-3-en-2-one) with

(sulfonylethane) using a catalyst (acetic anhydride) to obtain a heterogeneous cyclic compound with 6-membered ring (I) (Fig. 2).




Fig. 2. Synthesis of compound (I)

Using spectrometers (FT-IR, ^1H -NMR and ^{13}C -NMR), the nature of the prepared compound was confirmed.

The FT-IR of compound (I) shows a double stretching absorption band of the SO_2 group at 1420 and 1032 cm^{-1} . The appearance of a band at the range 3065 cm^{-1} is due to the $(=\text{C}-\text{H})$ group, a band at the range 2450 and 725 cm^{-1} is $(\text{C}-\text{H})$, and at the range 2655 cm^{-1} is due to the $(-\text{C}-\text{H})$ group $-\text{CH}_3$, and the functional group $(\text{C}=\text{C})$ appears at the range 1654 cm^{-1} and the appearance of a stretching band for the $(\text{C}-\text{O})$ group at the range 1266 cm^{-1} . The absence of a stretching band for the $(\text{C}=\text{O})$ group at the range 1740-1720 cm^{-1} .

The ^1H -NMR spectrum of compound (I) gave a signal for the CH_3-CO group at (2.81 s) δ ppm, and for $(\text{CH}_3)_2-\text{C}-$ at (2.05, 2.14 s) δ ppm, while the olefinic proton $\text{C}=\text{C}-\text{H}$ showed a signal at (6.91s) δ ppm, and a signal at (6.01-6.11 m) δ ppm. is due to the $\text{S}-\text{C}-\text{H}$ group.

^{13}C -NMR spectrum of compound (I) and chemical shifts $\text{d}_6\text{-DMSO}$. The appearance of the $\text{CH}_3-\text{C}-\text{SO}_2$ frequency at δ 19.98 ppm, and the methyl groups show two carbon atoms $(\text{CH}_3)_2-\text{C}-$

frequencies at 39.37-39.80 and 40.20-40.62 δ ppm, while the carbon atoms $=\text{C}-\text{CH}_3$ appear at 21.37 δ ppm. The frequency at 100.24 δ ppm, belongs to the carbon $(\text{CH}_3)_2-\text{C}-$ as well as the frequency of 113.96 δ ppm for the carbon $\text{CH}_3-\text{C}-\text{SO}_2$ and also the appearance of a frequency at position 147.09 δ ppm belongs to the carbon $\text{CH}_3-\text{C}=\text{C}-$ and also the frequency at 156.71 belongs to the carbon $\text{CH}_3-\text{C}=\text{C}$.

One of the most important reactions that these compounds (thiazines) undergo is the replacement of the ether oxygen atom in the 6-membered ring with the substituted nitrogen atom, via the ANRORC (nucleophilic ring addition-opening-ring-closing) reaction to produce it.

Then, to produce the last compound (II), 3 moles of 3,4,4,6-tetramethyl-3,4-dihydro-1,2-oxathiine 2,2-dioxide (I) react with one mole of benzene-1,3,5-triamine to obtain 2-(3,5-bis(3,5,5,6-tetramethyl-1,1-dioxido-5,6-dihydro-2H-1,2-thiazin-2-yl)phenyl)-3,5,5,6-tetramethyl-5,6-dihydro-2H-1,2-thiazine 1,1-dioxide (II) (Fig. 2).




Fig. 3. Synthesis of compound (II)

The FT-IR of compound (II) shows a double stretching absorption band for the SO_2 group at 1431 and 1023 cm^{-1} . A two bands at the

range 2966 cm^{-1} and 750 cm^{-1} are due to $(\text{C}-\text{H})$; the appearance of a band at the range 3063 cm^{-1} is due to the $(=\text{C}-\text{H})$ group; a band at the range

2961 cm^{-1} is due to the ($-\text{C}-\text{H}$) to $-\text{CH}_3$ group; and the functional group ($\text{C}=\text{C}$) appears at the range 1645 cm^{-1} , as well as two bands at positions 1317 and 1243 cm^{-1} due to the ($\text{C}-\text{N}$) group, and the absence of a stretching band for the ($\text{C}-\text{O}$) group at the range 1266-1420 cm^{-1} , as well as the disappearance of a double band due to ($-\text{NH}_2$) between the range 3000-3300 cm^{-1} .

The ^1H -NMR spectrum of compound (II) gave signals for the CH_3-CN group at (1.98 s) δ ppm and for $(\text{CH}_3)_2\text{C}=\text{C}$ at (5.01, 5.71 s) δ ppm, while the olefinic proton $\text{C}=\text{C}-\text{H}$ showed a signal at (7.54 s) δ ppm. Also, a signal at (7.31–7.42 M) δ ppm is due to the $\text{S}-\text{C}-\text{H}$ group, and a signal (7.58 s) δ ppm is due to the protons of the aromatic ring.

^{13}C -NMR spectrum of compound (II) and characteristic chemical shifts (d6- DMSO). The appearance of the $\text{CH}_3-\text{C}-\text{SO}_2$ frequency at 21.65 δ ppm, and the methyl groups show two carbon atoms $(\text{CH}_3)_2\text{C}-$ frequencies at 77.74, 77.36 δ ppm, while the carbon atoms $=\text{C}-\text{CH}_3$ appear at 29.37 δ ppm. as well as the frequency of 110.67 δ ppm for the carbon $\text{CH}_3-\text{C}-\text{SO}_2$ and also the appearance of a frequency at positions 129.54-129.74 and 129.86-129.64 δ ppm, which belongs to the olefinic carbon $\text{CH}_3-\text{C}=\text{C}-$; also the frequency at 141.61 δ ppm which belongs to the cyclo-olefinic carbon $\text{CH}_3-\text{C}-\text{N}-$; the frequency at 116.18 and 144.24 δ ppm is due to the aromatic ring carbons $\text{C}=\text{C}-\text{C}$ and $\text{C}=\text{C}-\text{N}$, respectively.




Fig. 4. Mechanism of synthesis of the two compounds (I) and (II)

Bioactivity. Using the cup agar plate diffusion method [39-42], the new synthesized compounds of 3,4,4,6-tetramethyl-3,4-dihydro-1,2-oxathine-2,2-dioxide (I) and 2-(3,5-bis(3,5,5,6-tetramethyl-1,1-dioxido-5,6-dihydro-2H-1,2-thiazin-2-yl)phenyl)-3,5,5,6-tetramethyl-5,6-dihydro-2H-1,2-thiazine-1,1-dioxide (II) were evaluated at concentrations (0.01, 0.1, 1.0, and 10 mg/ml) in the growth of six types of Gram-negative and positive bacteria (*Protans vulgarisginosa*, *Staphylococcus aureus*, *Pseudonas aeruginosa*, *Bacillus subtilus*, and *Enterbacter sp.*), using the drug compound (Chloramphenicol) as a reference. The inhibition

of compound (II) is observed to be greater than that of compound (I), in general, and for almost all concentrations. This is due to the increase in the number of faction groups when comparing the first and second synthesized compounds. A Table 1 shows high inhibition values at concentrations (1.0 and 10 mg/ml) and also exceeds the inhibition limit of the reference. It is also noted that the concentrations of the acid compound (0.01 and 0.1 mg/ml) are acceptable. This is done by measuring the inhibition values of the tablets (saturated with concentrations of the compound II) and embedded in bacteria dishes for 24 hours at a temperature of 37°C.

Table 1. The Diameter of Inhibition zone (mm)of Some Synthesized Compound Against Some Gram^{+ve} and Gram^{-ve} Bacteria

Comp. No.	Con. (mg/ml)	<i>Protans vulgarisginosa</i>	<i>Staphylo coccus aureus</i>	<i>Escheri chia coli</i>	<i>Bacillus subtilus</i>	<i>Enterbacter sp.</i>	<i>Pseudon as aerugin osa</i>
I	0.01	-	-	8	4	-	12
	0.1	4	-	9	12	5	13
	1.0	-	15	10	-	14	9
	10	10	9	11	20	17	15
II	0.01	2	14	-	10	24	14
	0.1	6	-	20	15	12	17
	1.0	10	27	19	16	25	24
	10	27	18	15	24	27	23
Chloramphenico l (30/μgm)disk	control	22	21	25	26	23	22

By observing the Table 2, which shows the effect of concentrations (10, 25, and 30 mg/ml) respectively for the synthetic compounds (I) and (II) against the types of fungi *A. fumigates*, *T.mentagrophytes*, *C.albicans* and *M. canis*. by measuring the inhibition zone (killing) for the concentrations of the prepared compounds, the killing range of compound (II) is much higher

compared to compound (I), due to the previous reason, which is the presence of a number of functional groups for compound (II) more than compound (I), and it is noted for all types of fungi with a very good evaluation except against *A. fumigates*, as it is noted that its resistance to killing is high.

Table 2. The diameter of inhibition zone (mm) for organic compounds at (10, 20, and 40 mg/ml) against some fungal species [43]

Fungal types	Diameter of inhibition zone (mm) for synthesizes of organic comp.					
	Cons. comp. (I)			Cons. comp. (II)		
	10	25	30	10	25	30
<i>T.mentagrophytes</i>	21	22	21	22	21	21
<i>C.albicans</i>	30	28	31	34	33	30
<i>A. fumigates</i>	16	13	14	17	15	10
<i>M. canis</i>	25	27	24	28	29	25

Molecular docking. Molecular docking studies were carried out for compounds (I) and (II) to identify the target enzymes under investigation and to assess the possible interactions of the heterocyclic rings, which have chemical affinities, indicating possible similarities in their chemical properties or methods, and amino acids as potential sources of importance in ligand binding. Docking studies of the present acidic residues ARG158 is an experimental compound with potential applications in medicine. Preliminary studies suggest that ARG158 may target biological pathways specifically associated with diseases [44-45]. Basic enzyme receptor studies influence

how it works, making it promising for further clinical research. Current research focuses on evaluating its efficacy, safety, and outcomes [46-48].

The study showed that compound (I) interacts with the protective residue at the site by forming various interactions: the hydrogen bonding ARG158 protective tank is located at the formation site with an oxygen pair of electrons for the sulfate complex in the vessel, and three alkyl bonds remain on security beads ILE254, MET160, and PRO98 located at the site of production with methyl groups in the mixture (Table 2, Fig. 5).




Fig. 5. The interaction between the compound (I) and Gastric cancer

Molecular docking studies of the prepared organic derivatives revealed the quantitative and qualitative interactions with amino acid residues in the active site, as the study identified compound (II) interacting with double bonds of amino acid residues ARG158, linking the sulfate

group of the compound to the two electrons of the oxygen atom, and double alkyl bonds linking amino acid residues in the active site with methyl groups of the compound LEU264 [49-51] (Tab. 3, Fig. 6).




Fig. 6. The interaction between the compound II and Gastric cancer

Table 3. Values of binding energies for the compound I and II

Comp. Symb.	RMSD	Docking Score
I	0.042	-7.7
II	0.046	-7.1

Conclusion

High-value heterocyclic compounds were synthesized for target compounds (I) and (II). Purified compounds were studied for effects on various bacteria (Gram-negative and Gram-positive) and were determined using spectroscopic techniques such as FT-IR, ^{13}C NMR, and ^1H -NMR, and structures were analyzed and discussed. The biological interactions of the formulations were compared

with those of conventional antibiotics. Our new compounds clearly showed inhibition of a variety of bacteria, as well as effects on fungi, some of which showed higher biological activity than the selected compounds in the two studies. A molecular docking study revealed several interactions involving amino acid residues in the active site.

References

1. Mondal S. Recent developments in the synthesis and application of sultones. *Chemical Reviews*, 2012, **Vol. 112(10)**, p. 5339-5355. DOI: [10.1021/cr2003294](https://doi.org/10.1021/cr2003294)
2. De Azevedo O.D.C.C., Heron B.M., Zonidis D. Synthesis, Reactivity and Applications of 1, 2-Oxathiine 2, 2-Dioxides. In *Targets in Heterocyclic Systems: Chemistry and Properties*, 2021, p. 33-56. DOI: 10.17374/targets.2021.24.33
3. Gaunersdorfer C., Waser M. Progress in the synthesis of δ -sultones. *Monatshefte für Chemie-Chemical Monthly*, 2018, **Vol. 149**, p. 701-714. DOI: 10.1007/s00706-017-2010-0

4. Evans D.A., Nagorny P., Rae K.J., Sonntag L.S., Reynolds D.J., Vounatsos F. Enantioselective synthesis of oasomycin A, Part III: Fragment assembly and confirmation of structure. *Angewandte Chemie*, 2007, **Vol. 119(4)**, p. 551-554. DOI: [10.1002/anie.200603652](https://doi.org/10.1002/anie.200603652)
5. Bequette J.P., Jungong C.S., Novikov A.V. Enantioselective synthesis of bakuchiol using diazosulfonate C–H insertion to install the quaternary center. *Tetrahedron letters*, 2009, **Vol. 50(50)**, p. 6963-6964. DOI: [10.1016/j.tetlet.2009.09.147](https://doi.org/10.1016/j.tetlet.2009.09.147)
6. Hamdoon A.M., Saleh M.Y. Synthesis & biological evaluation of novel series of benzo [f] indazole derivatives. *Egyptian Journal of Chemistry*, 2022, **Vol. 65(11)**, p. 305-312. DOI: [10.21608/EJCHEM.2022.120818.5418](https://doi.org/10.21608/EJCHEM.2022.120818.5418)
7. Farhan M.M., Rabeea M.A., Muslim R.F., Zidan T.A. Chemical composition (saturate fraction) of western Iraq natural bitumen. *Materials Today: Proceedings*, **Vol. 42**, p. 2527–2533. DOI: [10.1016/j.matpr.2020.12.574](https://doi.org/10.1016/j.matpr.2020.12.574)
8. Farhan M.M., Farhan F.M., Hussien A.M., Rahem H.J., Abdulmohsin I.M. Synthesis of a new compound to reduce the evaporation of petroleum and gasoline to protect the environment from pollution. In *AIP Conference Proceedings*. 2022, **Vol. 2394(1)**, 040026. DOI: [10.1063/5.0122217](https://doi.org/10.1063/5.0122217)
9. Al-Sultan R.M.H., Al-Sultan A.A., Hayawi M.A., Aldahham B.J., Saleh M.Y., Mohammed H.A. The effect of subclinical thyroid dysfunction on B-type natriuretic peptide level. *Revis Bionatura*, 2022, **Vol. 7(2)**, 21. DOI: [10.21931/RB/2022.07.02.21](https://doi.org/10.21931/RB/2022.07.02.21)
10. Farhan M.M., Najem R.A., Mandeel T.A.-J. Preparation of new organic compounds and the study of its characteristics to reduce the evaporation rate from gasoline tanks. *AIP Conference Proceedings*, 2020, **Vol. 2213**, 020216. DOI: [10.1063/5.0000354](https://doi.org/10.1063/5.0000354)
11. Hussein A.N., Fawzi M., Obaid R.F., Banoon S.R., Abood E.S., Ghasemian A. Dual pharmacological targeting of Mycobacterium tuberculosis (Mtb) PKNA/PKNB: A novel approach for the selective treatment of TB illness. *Egyptian Journal of Chemistry*, 2022, **Vol. 65(12)**, p. 409-419. DOI: [10.21608/EJCHEM.2022.135419.5981](https://doi.org/10.21608/EJCHEM.2022.135419.5981)
12. Moku B., Fang W.Y., Leng J., Li L., Zha G.F., Rakesh K.P., Qin H.L. Rh-catalyzed highly enantioselective synthesis of aliphatic sulfonyl fluorides. *Iscience*, 2019, **Vol. 21**, p. 695-705. DOI: [10.1016/j.isci.2019.10.051](https://doi.org/10.1016/j.isci.2019.10.051)
13. Zeng Y.Z., Wang J.B., Qin H.L. A reductive dehalogenative process for chemo- and stereoselective synthesis of 1, 3-dienylsulfonyl fluorides. *Organic & Biomolecular Chemistry*, 2022, **Vol. 20(39)**, p. 7776-7780. DOI: [10.1039/D2OB01434C](https://doi.org/10.1039/D2OB01434C)
14. Liu H., Moku B., Li F., Ran J., Han J., Long S., Qin H. L. Stereoselective Construction of Nitrile-Substituted Cyclopropanes from 2-Substituted Ethenesulfonyl Fluorides via Carbon-Sulfur Bond Cleavage. *Advanced Synthesis & Catalysis*, 2019, **Vol 361(19)**, p. 4596-4601. DOI: [10.1002/adsc.201900635](https://doi.org/10.1002/adsc.201900635)
15. Chen X., Zha G.F., Fang W.Y., Rakesh K.P., Qin H.L. A portal to a class of novel sultone-functionalized pyridines via an annulative SuFEx process employing earth abundant nickel catalysts. *Chemical Communications*, 2018, **Vol. 54(65)**, p. 9011-9014. DOI: [10.1039/C8CC04032J](https://doi.org/10.1039/C8CC04032J)
16. Huang Y.M., Wang S.M., Leng J., Moku B., Zhao C., Alharbi N.S., Qin H.L. Converting (E)-(Hetero) aryethanesulfonyl Fluorides to (Z)-(Hetero) aryethanesulfonyl Fluorides Under Light Irradiation. *European Journal of Organic Chemistry*, 2019, **Vol. 209(28)**, p. 4597-4603. DOI: [10.1002/ejoc.201900799](https://doi.org/10.1002/ejoc.201900799)
17. Aiken S., Gabbutt C.D., Heron B.M., Rice C.R., Zonidis D. Synthesis and photochromism of some mono and bis (thienyl) substituted oxathiine 2, 2-dioxides. *Organic & Biomolecular Chemistry*. 2019, **Vol. 17(44)**, p. 9578-9584. DOI: [10.1039/C9OB02128K](https://doi.org/10.1039/C9OB02128K)
18. Alrashedi H.S., Al-Ataie S.S., Banoon S.R., Fayed M.I. Potential role of medicinal plants for the treatment of respiratory viruses: A review. *Egyptian Journal of Chemistry*. 2021, **Vol. 64(12)**, p. 7495-7508. DOI: [10.21608/EJCHEM.2021.97912.4569](https://doi.org/10.21608/EJCHEM.2021.97912.4569)
19. Saied S.M., Saleh M.Y., Hamdoon A.M. Multicomponent synthesis of Tetrahydrobenzo [a] xanthene and Tetrahydrobenzo [a] acridine derivatives

- using Sulfonated Multi-Walled Carbon Nanotubes as Heterogeneous Nanocatalysts. *Iranian Journal of Catalysis*, (2022). 12(2), 189-205. DOI:10.30495/ijc.2022.1955651.1924
20. Farhan M.M., Younis A.A. Synthesis of new compounds of (5E, 5'E)-5, 5'-(1, 1'-biphenyl)-4, 4'-diylbis (azanylylidene)) bis (3, 3-dimethylcyclohexan-1-one) in new laboratory methods. *Egyptian Journal of Chemistry*, 2022, **Vol. 65(10)**, p. 515-522.
 21. Aiken S., Anozie K., Azevedo O.D., Cowen L., Edgar R.J., Gabbutt C.D., Zonidis D. Expedient synthesis of highly substituted 3, 4-dihydro-1, 2-oxathiine 2, 2-dioxides and 1, 2-oxathiine 2, 2-dioxides: revisiting sulfene additions to enaminketones. *Organic & Biomolecular Chemistry*, 2019, **Vol. 17(44)**, p. 9585-9604. DOI: [10.1039/C9OB01657K](https://doi.org/10.1039/C9OB01657K)
 22. Li B., Yan W., Zhang C., Zhang Y., Liang M., Chu F., Lei H. New synthesis method for sultone derivatives: Synthesis, crystal structure and biological evaluation of S-CA. *Molecules*, 2015, **Vol. 20(3)**, p. 4307-4318. DOI: [10.3390/molecules20034307](https://doi.org/10.3390/molecules20034307)
 23. Farhan M.M., Hussein M.A., Al-Jumaili M.H.A. Synthesis, characterization, and biological activities investigation of tri-armed structure of (7-bromo-5-chloro-2, 3-dihydrophthalazine-1, 4-dione) based on benzene ring. *Journal of Chemical Research*, 2024, **Vol. 48(6)**, DOI: [10.1177/17475198241306505](https://doi.org/10.1177/17475198241306505)
 24. Al-Saady A.J., Aldujaili N.H., Banoon S.R., Al-Abboodi A. Antimicrobial properties of nanoparticles in biofilms. *Revis Bionatura*. 2022, **Vol. 7(4)**, 71. DOI:10.21931/RB/2022.07.04.71
 25. Roudsari S.T., Rad-Moghadam K. A sulfonating ionic liquid for one-pot pseudo four-component synthesis of novel 3-chlorosulfonyl- δ -sultones: a novel class of fluorescent compounds. *Tetrahedron*, 2018, **Vol. 74(30)**, p. 4047-4052. DOI: [10.1016/j.tet.2018.06.012](https://doi.org/10.1016/j.tet.2018.06.012)
 26. Moon S., Khadtare S., Wong M., Han S.H., Bazan G.C., Choi H. Hole transport layer based on conjugated polyelectrolytes for polymer solar cells. *Journal of colloid and interface science*, 2018, **Vol. 518**, p. 21-26. DOI: [10.1016/j.jcis.2018.02.009](https://doi.org/10.1016/j.jcis.2018.02.009)
 27. Farhan M.M., Al-Jumaily M.M., Al-Muhammadi A.D., Ismail A.S. Development of a New Method for Reducing the Loss of Light Hydrocarbons at Breather Valve of Oil Tanks. *Energy Procedia*, 2017, **Vol. 141**, p. 471-478. DOI: [10.1016/j.egypro.2017.11.061](https://doi.org/10.1016/j.egypro.2017.11.061)
 28. Hussien B.M., Rabeea M.A., Farhan M.M. Characterization and behavior of Hydrogen Sulfide plumes released from active sulfidatar springs, Hit-Iraq. *Atmospheric Pollution Research*, 2020, **Vol. 11(5)**, p. 894-902. DOI: [10.1016/j.apr.2020.02.001](https://doi.org/10.1016/j.apr.2020.02.001)
 29. Hamdoon A.M., Al-Iraqi M.A., Saleh M.Y. Synthesis of some multi-cyclic sulfhydryl donor compounds containing 1, 2-dithiol-3-thione moiety. *Egyptian Journal of Chemistry*, 2022, **Vol. 65(3)**, p. 427-434. DOI: [10.21608/EJCHEM.2021.93344.4408](https://doi.org/10.21608/EJCHEM.2021.93344.4408)
 30. All-Heety M.M. Synthesis of New Compounds Bis-(N-Naphthal) Phthalisoimid, Bis-(N-Phenyl) Phthalisoimid and Studying Thermal Stability and Bio Activity. *Journal of Education and Science*. 2009, **Vol. 22(2)**, p.19-28. DOI: [10.33899/edusj.2009.57682](https://doi.org/10.33899/edusj.2009.57682)
 31. Qamar M.T., Sanz-Jimenez P., Banoon S.R., Zhu X.T., Aldakheel F.M., Almansour N.M., Ahmad F. Computational identification of anti-cancer compounds targeting the RNA-binding domain of human FOX-1 protein (RBFOX1). *Results in Chemistry*, 2025, **Vol. 13**, 102004. DOI: [10.1016/j.rechem.2024.102004](https://doi.org/10.1016/j.rechem.2024.102004)
 32. Hassan B.A., Lawi Z.K.K., Banoon S.R. Detecting the activity of silver nanoparticles, pseudomonas fluorescens and Bacillus circulans on inhibition of aspergillus Niger growth isolated from moldy orange fruits. *Periódico Tchê Química*, 2020, **Vol. 17(35)**, p. 678-690. DOI: [10.52571/PTQ.v17.n35.2020.58](https://doi.org/10.52571/PTQ.v17.n35.2020.58) [BANOON_pgs_678_690.pdf](#)
 33. Mahmood Z.M., Saied S.M., Saleh M.Y. Deep eutectic solvent (reline) as catalytic system in synthesis of schiff bases derived from glucose and different amines. *Chemical Problems*, 2024, **Vol. 22(3)**, p. 281-289. DOI: [10.32737/2221-8688-2024-3-281-289](https://doi.org/10.32737/2221-8688-2024-3-281-289)
 34. Ali A.H., Saleh M.Y., Yaqoob Q.A., Saied S.M., Hasan M.S., Owaid K.A., Alsirhani

- A.M. Comprehensive evaluation of antibacterial and anticancer activities from indole butanoic acid. *Journal of Genetic Engineering and Biotechnology*, 2025, **Vol. 23(1)**, 100452. DOI: [10.1016/j.jgeb.2024.100452](https://doi.org/10.1016/j.jgeb.2024.100452)
35. Ali R.T., Al-Niemi K.I., Mohammedthalji N.H. A practical and theoretical study of the mechanical kinetics of ascorbic acid adsorption on a new clay surface. *Egyptian Journal of Chemistry*, 2021, **Vol. 64(8)**, p. 4569-4581. DOI: [10.21608/EJCHEM.2021.71632.3572](https://doi.org/10.21608/EJCHEM.2021.71632.3572)
36. Doss S. H., Baghos V. B., Abdelhamid A. O., Halim M. M. A. Synthesis of Sultam derivatives with expected biological activity. *Molecules*, 2000, **Vol. 5(6)**, p. 816-825. DOI: [10.3390/50600816](https://doi.org/10.3390/50600816)
37. Tornøe C.W., Christensen C., Meldal M. Peptidotriazoles on solid phase:[1, 2, 3]-triazoles by regiospecific copper (I)-catalyzed 1, 3-dipolar cycloadditions of terminal alkynes to azides. *The Journal of organic chemistry*, 2002, **Vol. 67(9)**, p. 3057-3064. DOI: [10.1021/jo011148j](https://doi.org/10.1021/jo011148j)
38. Silverstein R.M., Bassler G.C. Spectrometric identification of organic compounds. *Journal of Chemical Education*, 1962, **Vol. 39(11)**, 546. DOI: [10.1021/ed039p546](https://doi.org/10.1021/ed039p546)
39. Victoria D.B., Henk Hiemstra J.H. Cu^I-Catalyzed Alkyne-Azide "Click" Cycloadditions from a Mechanistic and Synthetic Perspective. *European Journal Organic Chemistry*. 2006, **Vol. 55(1)**, p. 51-68. DOI: [10.1002/ejoc.200500483](https://doi.org/10.1002/ejoc.200500483)
40. Wu P., Feldman A.K., Nugent A.K., Hawker C.J., Scheel A., Voit B., Pyun J., Fréchet J.M.J., Sharpless K.B., Fokin V.V. Efficiency and Fidelity in a Click-Chemistry Route to Triazole Dendrimers by the Copper(I)-Catalyzed Ligation of Azides and Alkynes. *Angew. Chem. Int. Ed*, 2004, **Vol. 43**, p. 3928-3932. DOI: [10.1002/anie.200454078](https://doi.org/10.1002/anie.200454078)
41. Farhan M.M., Al-Jumaili M.H.A., Bakr E.A. New bis-heterocyclic structures: Synthesis, characterization and biological activity. *Journal of Chemical Research*, 2023, **Vol. 47(6)**, 17475198231218387. DOI: [10.1177/17475198231218387](https://doi.org/10.1177/17475198231218387)
42. Banoon Z.R., Mahmood R.S., Hamad A.R., Hussein Z.A. Design, microwave synthesis, characterization and antimicrobial activity of imidazolone derivatives. *Journal of Molecular Structure*, 2025, **Vol. 1322(4)**, 140701. DOI: [10.1016/j.molstruc.2024.140701](https://doi.org/10.1016/j.molstruc.2024.140701)
43. Farhan M.M. A comparative study of oxidation in the preparation of lacton and bislacton compounds and biological activity study of product compounds. *Journal of Global Pharma Technology*, 2009, **Vol. 10(05)**, p. 484-496
44. Sadeek G.T., Saeed Z.F., Saleh M.Y. Synthesis and pharmacological profile of hydrazide compounds. *Research Journal of Pharmacy and Technology*, 2023, **Vol. 16(2)**, p. 975-982. DOI: [10.52711/0974-360X.2023.00163](https://doi.org/10.52711/0974-360X.2023.00163)
45. Vandepitte J., Engback K., Piot P., Heuck C.C. Basic microbiology procedures in clinical bacteriology. *World Health Organisation, Geneva*. 1991, 2 nd ed. 167 p.
46. Saleh M.Y., Aldulaimi A.K.O., Saeed S.M., Adhab A.H. TiFe₂O₄@ SiO₂-SO₃H: A novel and effective catalyst for esterification reaction. *Heliyon*, 2024, **Vol. 10(4)**, p. 546-357. DOI: [10.1016/j.heliyon.2024.e26286](https://doi.org/10.1016/j.heliyon.2024.e26286)
47. Saleh M.Y., Aldulaimi A.K.O., Saeed S.M., Adhab A.H. Palladium fabricated on Fe₃O₄ as an organic-inorganic hybrid nanocatalyst for the Suzuki and Stille coupling reactions. *Journal of Molecular Structure*, 2025, **Vol. 132(11)**, p. 1395-1397. DOI: [10.1016/j.molstruc.2024.139597](https://doi.org/10.1016/j.molstruc.2024.139597)
48. Farhan M.M., Guma M.A., Rabeea M.A., Ahmad I., Patel H. Synthesizes, characterization, molecular docking and in vitro bioactivity study of new compounds containing triple beta lactam rings. *Journal of Molecular Structure*, 2022, **Vol. 1269**, 133781. DOI: [10.1016/j.molstruc.2022.133781](https://doi.org/10.1016/j.molstruc.2022.133781)
49. Ali A.H., Saleh M.Y., Owaid K.A. Mild Synthesis, Characterization, and Application of some Polythioester Polymers Catalyzed by Cetrinide Ionic Liquid as a Green and Eco-Friendly Phase-Transfer Catalyst. *Iranian Journal of Catalysis*. 2023, **Vol. 13(1)**, p. 36-45. DOI: [10.30495/ijc.2023.1973500.1977](https://doi.org/10.30495/ijc.2023.1973500.1977)

-
50. Al-Jubori H.M.S., Al-Mathkuri T.S.F., Banoon Z.R., Saleh M.Y. Synthesis of novel benzo[d] imidazole bearing α -aminophosphonate and their antimicrobial evaluation. *Results in Chemistry*, 2024, **Vol. 8**, 101586. DOI: [10.1016/j.rechem.2024.101586](https://doi.org/10.1016/j.rechem.2024.101586)
51. Ahmad H.M., Aldahham B.J., Saleh M.Y. Dehydroepiandrosterone supplementation improves diminished ovarian reserve clinical and in silico studies. *Steroids*. 2024, **Vol. 211(3)**, p. 109-115. DOI: [10.1016/j.steroids.2024.109490](https://doi.org/10.1016/j.steroids.2024.109490)