

PREPARATION, BIOLOGICAL EVALUATION, AND APPLICATION OF THIADIAZOLE, SCHIFF BASE, AND TETRAZOLINE DERIVATIVES DERIVED FROM AQUEOUS MORINGA LEAF EXTRACTS: POM ANALYSIS FOR PHARMACOPHORE SITE IDENTIFICATION

Hadil M. Salih^{1,2}, Sanaa A. Abd-Alkareem¹

¹Department of Chemistry, College of Science for Woman, University of Baghdad, Iraq

²Scientific Research Commission, Baghdad, Iraq

e-mail: Hadeel.Mahdi2305m@cs.w.uobaghdad.edu.iq, sanaaa_chem@cs.w.uobaghdad.edu.iq

Received 31.08.2025

Accepted 17.11.2025

Abstract: *Moringa oleifera* leaves have long been used in conventional medicine to treat a variety of ailments, including fever and sore throat. The close relationship between plants and human health has promoted the development of plant-based therapies, such as pharmaceuticals and nutritional supplements, thereby advancing traditional medicine and adding value to agriculture. In the present study, novel thiadiazole, Schiff base, and tetrazoline derivatives (H1, H2, and H3) were synthesized from nitrile-containing precursors obtained from the aqueous extract of *Moringa oleifera* leaves. The synthesis involved the reaction of the nitrile compound with thiourea in the presence of glacial acetic acid to form a heterocyclic thiadiazole intermediate, followed by condensation with an aromatic aldehyde to yield Schiff bases, and subsequent reaction with sodium azide to produce tetrazoline derivatives.

The synthesized compounds were characterized by Fourier-transform infrared spectroscopy (FT-IR), melting point determination, and ¹H NMR spectroscopy. The biological activities of derivatives H1–H3 were evaluated against *Staphylococcus aureus* (Gram-positive), *Escherichia coli* (Gram-negative), and the fungal strain *Candida albicans*. Compared with standard drugs—ciprofloxacin for antibacterial activity and ketoconazole for antifungal activity—compound H3 exhibited superior antibacterial and antifungal activities. In addition, H3 demonstrated excellent antioxidant activity relative to ascorbic acid. POM analysis was further employed to identify promising pharmacophore sites within the synthesized compounds, highlighting their potential for pharmaceutical applications.

Keywords: Nitril Compounds, thiourea, Schiff bases, cyclic addition reaction, tetrazoline, heterocyclic.

DOI: 10.65382/2221-8688-2027-1-181-191

Introduction

Moringa oleifera is an important medicinal plant rich in vitamins, minerals, flavonoids, and amino acids [1]. It belongs to the family Moringaceae and is a fast-growing tree that can reach heights of up to 10 m, characterized by its abundant green leaves [2]. Due to its wide availability and diverse phytochemical composition, *Moringa oleifera* has attracted considerable attention in medicinal and pharmaceutical research.

One of the primary objectives of organic synthesis is to develop efficient and sustainable synthetic routes to biologically active compounds using readily available starting materials. Among heterocyclic systems, 1,3,4-thiadiazole represents a five-membered

heterocyclic scaffold with distinctive physicochemical properties. Its derivatives are well known for exhibiting a broad spectrum of biological activities, including antibacterial [3] and anti-Alzheimer effects [4,5]. Moreover, the 1,3,4-thiadiazole ring is an integral structural motif in several clinically used drugs, such as acetazolamide, methazolamide, cefazolin, cefazedone, sulfamethizole, and megazol [6].

Schiff bases constitute another important class of organic compounds due to their diverse biological activities, including antimalarial [7], antitumor [8], antifungal [9], antibacterial [10], antitubercular [11], antiviral [12], anticancer [13], and antiproliferative properties [14]. These compounds are typically formed through the

condensation reaction of a primary amine with an aldehyde or ketone, often facilitated by catalysts such as glacial acetic acid. Structurally, Schiff bases are characterized by the presence of the azomethine ($-C=N-$) functional group, which plays a crucial role in their biological activity [15]. The field of synthetic organic chemistry continues to advance the development of bioactive compounds, innovative drug designs, and efficient synthetic methodologies [16, 17]. Heterocyclic compounds have attracted particular attention because they constitute the structural core of numerous synthetic and naturally occurring medicinal agents [18, 19]. Among these, nitrogen-containing heterocycles are especially significant in the life sciences due to their structural diversity, ability to engage in intermolecular and enzyme interactions, electronic versatility, natural abundance, and widespread use as key intermediates in medicinal chemistry for the development of novel pharmaceuticals [20, 21].

Tetrazoles are an important class of

heterocyclic compounds composed of a five-membered ring containing four nitrogen atoms and one carbon atom, and they can be obtained either by synthesis or from natural derivatives [22, 23]. The incorporation of the tetrazole moiety into bioactive molecules represents a rational strategy in drug design, as it can improve pharmacokinetic properties while reducing undesirable effects of potent drug candidates [24]. Cycloaddition reactions are among the most significant transformations in organic chemistry, owing to their synthetic utility and mechanistic importance [25].

In addition, by closely examining the effects of the most promising compounds on microbial cell membranes and evaluating their antioxidant properties, this study aims to gain deeper insight into their mechanisms of action. The findings are expected to contribute to the development of new and effective antimicrobial agents for the treatment of bacterial and fungal infections.

Experimental part

Materials. The chemicals used in this study were obtained from various commercial suppliers. Thiourea (99%) was purchased from Kem Aus, absolute ethanol (99%) from GCC, and glacial acetic acid, sodium azide, and aromatic amines were obtained from standard analytical suppliers. Dimethyl sulfoxide (DMSO, 99%) was supplied by BJH and used as the solvent.

Methods. Gas chromatography–mass spectrometry (GC–MS) analyses were performed using a PerkinElmer GC–MS system. Fourier-transform infrared (FT-IR) spectra were recorded on a Shimadzu 8400S spectrometer. Melting points were determined using a digital CG-1839-A0 melting point apparatus.

Synthesis of Thiadiazole Derivative 2-((2-(5-Amino-1,3,4-thiadiazol-2-yl)ethyl)(methyl)amino)-1-phenylpropan-1-ol (H1) [26]. The starting nitrile compound, 2-((1-hydroxy-1-phenylpropan-2-yl)(methyl)amino)acetonitrile (0.005 mol, 0.012 g), was dissolved in 25 mL of ethanol. Thiourea (0.010 mol, 0.74 g) was added to the reaction mixture, followed by the dropwise addition of glacial acetic acid. The resulting mixture was

refluxed for 12 h. After completion of the reaction, the mixture was cooled, filtered, and the crude product was recrystallized from ethanol to afford the thiadiazole derivative H1.

Synthesis of Schiff Base Derivative **(Z)-2-((2-(5-((4-Chlorobenzylidene)amino)-1,3,4-thiadiazol-2-yl)ethyl)(methyl)amino)-1-phenylpropan-1-ol (H2)** [27]. A mixture of thiadiazole compound H1 (0.001 mol, 0.292 g) and *p*-chlorobenzaldehyde (0.001 mol, 0.14 g) was dissolved in 20 mL of absolute ethanol. Several drops of glacial acetic acid were added as a catalyst, and the reaction mixture was refluxed for 6–8 h. Upon completion, the mixture was cooled and filtered, and the product was recrystallized from absolute ethanol to yield the Schiff base derivative H2.

Synthesis of Tetrazoline Derivative **Sodium 5-(4-chlorophenyl)-4-(5-(2-((1-hydroxy-1-phenylpropan-2-yl)(methyl)amino)ethyl)-1,3,4-thiadiazol-2-yl)-4,5-dihydrotetrazol-1-ide (H3)** [27]. The tetrazoline derivative (H3) was synthesized by reacting the Schiff base derivative H2 (0.001 mol, 0.414 g) with sodium azide (NaN_3 , 0.001 mol, 0.07 g) in dimethylformamide (DMF) as the

solvent. Copper(I) chloride (CuCl) was added as a catalyst. The reaction mixture was stirred at room temperature for approximately 24 h. Upon completion, the reaction mixture was filtered,


and the solid product was washed with distilled water, dried, and finally recrystallized from absolute ethanol to afford the tetrazoline compound H3.

Result and discussion

Due to their wide range of applications, particularly in pharmaceutical, industrial, and agricultural fields, heterocyclic compounds have attracted significant research interest. Accordingly, in this work, a series of heterocyclic derivatives—including thiadiazole, Schiff base, and tetrazoline compounds—were synthesized.

The thiadiazole derivative H1 was prepared through the reaction of a nitrile-containing compound, 2-((1-hydroxy-1-phenylpropan-2-yl)(methyl)amino)acetonitrile,

with thiourea. The Schiff base derivative H2, namely (Z)-2-((2-(5-((4-chlorobenzylidene)amino)-1,3,4-thiadiazol-2-yl)ethyl)(methyl)amino)-1-phenylpropan-1-ol, was synthesized by the condensation reaction of compound H1 with an aromatic aldehyde. Subsequently, the tetrazoline derivative H3 was obtained via cycloaddition of the Schiff base H2 with sodium azide in the presence of copper(I) chloride as a catalyst. The overall synthetic route is illustrated in Scheme 1.



Scheme 1. Synthetic steps for preparation of derivatives[H₁],[H₂],[H₃]

The structural formulas, yields (%), melting points (Mp), colors, and FT-IR spectral data of the synthesized compounds are summarized in Table 1. Among the prepared derivatives, compound H3 exhibited the highest melting point (>250 °C), whereas the lowest melting point was observed for compound H2 (140–142 °C). The highest reaction yield was obtained for compound H3, while compound H1 showed the lowest yield.

In the FT-IR spectra (Fig.s 1-3), the characteristic absorption bands of the amino group (NH₂) appeared in the range 3284–3278 cm⁻¹, and the $\nu(\text{C-S})$ stretching vibrations were observed at 630, 547, and 522 cm⁻¹. The Schiff base derivative H2 displayed a distinctive azomethine (C=N) stretching band at 1685 cm⁻¹ [28]. Other characteristic absorption bands are listed in Table 1.

Table 1. Physicochemical properties and FT-IR spectral data (cm⁻¹) of the synthesized compounds H1, H2, and H3

Comp.	Structure of components	Mp, (°C)	Color	Yield %	V(C-H) aromatic	V(C-H) Aliphatic	Other bands
-------	-------------------------	----------	-------	---------	-----------------	------------------	-------------

H1		165- 167	Thin white sheets	70	3105	2925-2856	NH ₂ :3384-3278 C-S:630 N-N(aromatic):1083 C=N:1467
H2		140-142	White powder	75	3095	2925-2852	N=CH:(imine)1685 C-Cl:761 C-S:547 N-N:(aromatic)1089
H3		248-250	Black powder	80	3130	2975-2925	C-S:522 C-Cl:775 N ₃ :2063 N=N(aromatic):1596




Fig. 1. FTIR spectrum of compound H1




Fig. 2. FTIR spectrum of compound H2




Fig. 3. FTIR spectrum of compound H3

The $^1\text{H-NMR}$ spectrum of compound H1 (Fig. 4) explain chemical shifts, δ (ppm) showed 5.45 (s, H, NH_2), 3.26 (s, H, OH), 1.24-1.51(d, 3H, $\text{CH}_3\text{-CH}$), 2.29 (s, 3H, $\text{CH}_3\text{-N}$) 2.74-2.93(t, 4H, $\text{CH}_2\text{-CH}_2$), 3.14-3.87 (t, 1H, CH-N), 6.22-5.45 (s, H, NH_2), 3.26 (s, H, OH), 1.24-1.51(d, 7.08 (m, 5H, Ar-H) [29].




Fig. 4. $^1\text{H-NMR}$ spectra of compound H1

The $^1\text{H-NMR}$ spectrum of compound H2 (Fig. 5) explain chemical shifts, δ (ppm) showed 3.64 (s, H, OH), 1.25-1.52 (d, 3H, $\text{CH}_3\text{-CH}$), 2.30 (s, 3H, $\text{CH}_3\text{-N}$) 2.32-2.90 (t, 4H, $\text{CH}_2\text{-CH}_2$), 3.13-3.33 (t, 1H, CH-N), 4.64-4.35 (d, 1H, CH-O), 6.78-7.72 (m, 9H, Ar-H), 9.68 (s, 1H, $\text{CH}=\text{N}$) [30].




Fig. 5. $^1\text{H-NMR}$ spectra of compound H2

The Biological Activity Studies [31-34]. The biological activity of newly synthesized pharmaceutical compounds—namely thiadiazole, Schiff base, and tetrazoline derivatives (H1, H2, and H3)—prepared from a nitrile compound using *Moringa oleifera* leaf extract was investigated. These compounds were selected as the main bioactive components for antimicrobial evaluation.

Biological assays were carried out *in vitro* against two bacterial strains, *Staphylococcus aureus* (Gram-positive) and *Escherichia coli* (Gram-negative), as well as one fungal strain,

Candida albicans. The antibacterial activity was evaluated using Mueller–Hinton agar, while antifungal activity was assessed using potato dextrose agar, due to their widespread application in medical microbiology.

Compound H1 exhibited moderate to weak antibacterial activity against the tested bacterial strains, whereas compound H3 showed a pronounced antibacterial effect, ranging from strong to moderate activity. The antibacterial activity data for all synthesized compounds are summarized in Table 2 and Fig. 6.

Table 2. Antibacterial activity of prepared compounds H1-H3

Compd.	<i>S.aureu</i>	<i>Escherichia coli</i>	<i>Candida albicans</i>
H1	13	11	15
H2	12	15	10
H3	27	13	15




Fig. 6. Biological activity for synthesized compounds H1-H3

Antioxidant Studies [35, 36]. In this study, the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay was employed as a simple, inexpensive, and widely used method to evaluate the antioxidant activity of free radical-scavenging compounds. This method assesses the ability of substances to act as hydrogen donors or free radical scavengers.


The antioxidant activity of *Moringa oleifera* extract derivatives (H1, H2, and H3) was evaluated *in vitro* using the DPPH method. At a concentration of 0.5 µg/mL, compound H2 exhibited significantly lower antioxidant activity, whereas compound H3 showed the

highest antioxidant activity among the tested compounds. Compound H1 displayed the lowest IC₅₀ value (17.26 µg/mL), indicating the strongest radical-scavenging activity, while compound H2 showed a higher IC₅₀ value (25.23 µg/mL). For comparison, the standard antioxidant ascorbic acid exhibited an IC₅₀ value of 33.3 µg/mL.

The antioxidant activity of all compounds decreased with decreasing concentration. The antioxidant activity and IC₅₀ values of the investigated compounds are summarized in Table 3.

Table 3. Antioxidants and IC₅₀ for the derivatives (H1, H2, H3)

Compd.	Con. (ppm)	I (%)	IC ₅₀
H1	0.5	79.522	17.27
	0.25	68.645	
	0.125	52.114	
	0.0625	40.025	
H2	0.5	77.541	25.23
	0.25	65.541	
	0.125	42.521	
	0.0625	35.521	
H3	0.5	85.542	22.78
	0.25	72.212	
	0.125	59.854	
	0.0625	44.541	
Ascorbic acid	0.5	83.07	33.78
	0.25	65.92	
	0.125		
	0.0625		

**Fig. 7.** The anti-oxidant studies for H1, H2 and H3

POM (PETRA/OSIRIS/Molinspiration) [33–35]. The toxicity risks and molecular properties of compounds H1–H3 were evaluated using the PETRA, OSIRIS, and Molinspiration computational programs. OSIRIS analysis indicated that all investigated compounds

Theory




possess favorable drug scores (DS = 63–93%) and are predicted to be non-mutagenic, non-tumorigenic, and non-toxic to reproductive systems.

Molinspiration analysis further revealed that compounds H1–H3 comply fully with Lipinski's rule of five, exhibit good drug-

likeness, and show no rule violations. The predicted pharmacokinetic properties and drug-likeness probabilities of these compounds are summarized in Table 4. POM analysis predicted that the physicochemical properties of the studied compounds enable effective antibacterial and antifungal activity, which is attributed to the

presence of pharmacophoric sites such as $\text{NH}^{\delta+}$ and $\text{S}^{\delta-}$ groups. Compounds H1–H3 contain combined pharmacophore features, and their biological activity is influenced by the spatial distances between polar atoms ($\text{NH}^{\delta+} \cdots \text{S}^{\delta-}$), as presented in Table 4.

Table 4. Osiris and molinspiration calculations of molecular properties of (H1-H3)

No. Compd.	Structures	Osiris calculations				Molinspiration calculations	
H1		Toxicity Risks	cLogP	TPSA		miLogP	1.22
		mutagenic	1.02	103.6		MW	292.41
		tumorigenic	Solubility	Druglikeness		nON	5
		irritant	-1.79	8.94		nOHNNH	3
		reproductive effective	Molweight	Drug-Score		nviolations	0
H2		Toxicity Risks	cLogP	TPSA		miLogP	3.95
		mutagenic	3.67	89.85		MW	414.96
		tumorigenic	Solubility	Druglikeness		nON	5
		irritant	-3.99	10.11		nOHNNH	1
		reproductive effective	Molweight	Drug-Score		nviolations	0
H3		Toxicity Risks	cLogP	TPSA		miLogP	0.57
		mutagenic	3.88	105.4		MW	479.0
		tumorigenic	Solubility	Druglikeness		nON	8
		irritant	-3.37	8.07		nOHNNH	1
		reproductive effective	Molweight	Drug-Score		nviolations	0
			479.0	0.63		volume	391.40




Fig. 10. Atomic Charge for components (H1-H3)

Conclusion

This study aimed to prepare an aqueous extract of *Moringa oleifera* leaves and to identify its active constituents under laboratory conditions. The most effective compounds were selected following investigation by gas chromatography–mass spectrometry (GC–MS). A nitrile compound was subsequently used as a precursor for the synthesis of thiadiazole, Schiff base, and tetrazoline derivatives (H1, H2, and


H3). The structures of the synthesized compounds were confirmed using proton nuclear magnetic resonance (^1H NMR) and Fourier transform infrared (FTIR) spectroscopy.

The newly synthesized compounds were evaluated for their *in vitro* antibacterial, antifungal, and antioxidant activities. Several derivatives exhibited promising biological activity, with compound H3 showing the most

pronounced antibacterial, antifungal, and antioxidant effects.

Moreover, a good correlation was observed between the experimental biological results and the computational studies based on POM analysis. The in silico evaluation indicated that the investigated heterocyclic compounds possess a favorable non-toxic profile, good

bioavailability, and acceptable pharmacokinetic properties. These encouraging results support further investigation, particularly focusing on the possible metabolic pathways of the thiadiazole ring under acidic physiological conditions. The proposed metabolite structure is illustrated in Equation 1:



Conflict of interest

The authors announced that there are no conflicts of interest regarding this current work.

References

- Islam Z., Islam S.M.R., Hossen F., Mahtab-Ul-Islam K., Hasan M.R., Karim R. Moringa oleifera is a Prominent Source of Nutrients with Potential Health Benefits. *International Journal of Food Science*, 2021, **Vol. 2021**, 6627265. DOI: [10.1155/2021/6627265](https://doi.org/10.1155/2021/6627265)
- Pareek A., Pant M., Gupta M.M., Kashania P., Ratan Y., Jain V., Pareek A., Chuturgoon A.A. Moringa oleifera: An Updated Comprehensive Review of Its Pharmacological Activities, Ethnomedicinal, Phytopharmaceutical Formulation, Clinical, Phytochemical, and Toxicological Aspects. *International Journal of Molecular Sciences*, 2023, **Vol. 24**, 2098. DOI: [10.3390/ijms24032098](https://doi.org/10.3390/ijms24032098)
- Anthwal T., Paliwal S., Nain S. Diverse Biological Activities of 1,3,4-Thiadiazole Scaffold. *Chemistry*, 2022, **Vol. 4**, p. 1654–1671. DOI: [10.3390/chemistry4040107](https://doi.org/10.3390/chemistry4040107)
- Bansal A.K., Antolini F., Zhang S., Stroea L., Ortolani L., Lanzi M., Serra E., Allard S., Scherf U., Samuel I.D.W. Highly luminescent colloidal CdS quantum dots with efficient near-infrared electroluminescence in light-emitting diodes. *Journal of Physical Chemistry C*, 2016, **Vol. 120**, p. 1871–1880. DOI: [10.1021/acs.jpcc.5b09109](https://doi.org/10.1021/acs.jpcc.5b09109)
- Indelicato S., Bongiorno D., Mauro M., Cascioferro S. Recent Developments of 1,3,4-Thiadiazole Compounds as Anticancer Agents. *Pharmaceuticals (Basel)*, 2025, **Vol. 18**, 580. DOI: [10.3390/ph18040580](https://doi.org/10.3390/ph18040580)
- Serban G. Synthetic Compounds with 2-Amino-1,3,4-Thiadiazole Moiety Against Viral Infections. *Molecules*, 2020, **Vol. 25**, 942. DOI: [10.3390/molecules25040942](https://doi.org/10.3390/molecules25040942)
- Kumar A., Verma S., Mishra A.K., Kumar S. Synthesis of some new Schiff bases of Pharmaceutical Interest. *Annals of Advanced Chemistry*, 2017, **Vol. 1**, p. 053–056. DOI: [10.29328/journal.aac.1001006](https://doi.org/10.29328/journal.aac.1001006)
- Shakeela Y., Sajjad H.S., Muhammad S.A., Zahid H.C. Antimicrobial metal-based thiophene derived compounds. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 2017, **Vol. 32**, p. 106–112. DOI: [10.1080/14756366.2016.1238363](https://doi.org/10.1080/14756366.2016.1238363)
- Suman M., Bharti J., Amar S.M., Archana S. Review Study on Metal Complexes Synthesized from Therapeutically Important Schiff Bases. *The Pharmaceutical and Chemical Journal*, 2016, **Vol. 3**, p. 119–124.
- Mabrouk M.S., Soad G.A., Safaa S.H. Synthesis, Characterizations, Biological, and Molecular Docking Studies of Some Amino Acid Schiff Bases with Their Cobalt(II) Complexes. *Advances in*

- Biological Chemistry*, 2017, **Vol. 7**, p. 182–194. DOI: [10.4236/abc.2017.75013](https://doi.org/10.4236/abc.2017.75013)
11. Wagdy M.E., Mohamed F., Marwa M.A., Hatem A.A. Design, Synthesis and Antitubercular Activity of Certain Nicotinic Acid Hydrazides. *Molecules*, 2015, **Vol. 20**, p. 8800–8815. DOI: [10.3390/molecules20058800](https://doi.org/10.3390/molecules20058800)
 12. Al-Azzawi A.M., Ali A.R. Synthesis and Antibacterial Screening of New Schiff Bases Based on N-(4-acetophenyl) Succinimide. *Journal of Science*, 2017, **Vol. 58**, p. 1790–1801. DOI: [10.24996/ijcs.2017.58.4A.2](https://doi.org/10.24996/ijcs.2017.58.4A.2)
 13. Miri R.R., Azzaghi-asl N., Mohammadi M.K. Study and conformational analysis of an isatin Schiff base as a potential cytotoxic agent. *Journal of Molecular Modeling*, 2013, **Vol. 19**, p. 727–735. DOI: [10.1007/s00894-012-1586-x](https://doi.org/10.1007/s00894-012-1586-x)
 14. Ashu C., Anshul S. Schiff bases: An emerging potent class of pharmaceuticals. *International Journal of Current Research in Medical Sciences*, 2017, **Vol. 3**, p. 60–74. DOI: [10.22192/ijcrms.2017.03.06.009](https://doi.org/10.22192/ijcrms.2017.03.06.009)
 15. Ajay K., Shweta V., Arun K.M., Sushil. Synthesis of some new Schiff bases of Pharmaceutical Interest. *Annals of Advanced Chemistry*, 2017, **Vol. 1**, p. 053–056. DOI: [10.29328/journal.aac.1001006](https://doi.org/10.29328/journal.aac.1001006)
 16. Alsahib S.A. Characterization and Biological Activity of Some New Derivatives Derived from Sulfamethoxazole Compound. *Baghdad Science Journal*, 2020, **Vol. 17**, p. 471–480. DOI: [10.21123/bsj.2020.17.2.0471](https://doi.org/10.21123/bsj.2020.17.2.0471)
 17. Hussein I.A., Narren S.F. Synthesis and characterization of some new derivatives starting from (4,4'-diaminophenoxy) ethane containing diazonium groups. *Journal of Global Pharma Technology*, 2019, **Vol. 11**, p. 97–107.
 18. Mezaal N.N., Alsahib S.A. Synthesis and characterization of new derivatives using Schiff's bases for alcoholic extract of (*Cordia myxa*) medical plant. *Eurasian Chemical Communications*, 2022, **Vol. 4**, p. 1054–1061. DOI: [10.22034/ecc.2022.343900.1473](https://doi.org/10.22034/ecc.2022.343900.1473)
 19. Narran S.F., Mohammed S.S., Omer M.K., Hussein I.A., Jawad N.M., Shweish B.K. Synthesis and Biological Activities of Some New Derivatives Based on 5-Styryl-2-amino-1,3,4-thiadiazole. *Chemical Methodologies*, 2022, **Vol. 6**, p. 83–90. DOI: [10.22034/chemm.2022.2.1](https://doi.org/10.22034/chemm.2022.2.1)
 20. Kerru N., Gummidi L., Maddila S., Gangu K.K., Jonnalagadda S.B. A Review on Recent Advances in Nitrogen-Containing Molecules and Their Biological Applications. *Molecules*, 2020, **Vol. 25**, 1909. DOI: [10.3390/molecules25081909](https://doi.org/10.3390/molecules25081909)
 21. Thejeel K., Abdul Kareem S., Ascar I., Hussein M. Synthesis of new polymers linked to heterocyclic using zinc oxide with nanostructures extracted from natural sources. *Egyptian Journal of Chemistry*, 2022, **Vol. 65**, p. 579–589. DOI: [10.21608/ejchem.2021.92971.4644](https://doi.org/10.21608/ejchem.2021.92971.4644)
 22. Abdulridha M.Q., Al-Hamdani A.A.S., Hussein I.A. Synthesis, Characterization and Antioxidant Activity of New Azo Ligand and Some Metal Complexes of Tryptamine Derivatives. *Baghdad Science Journal*, 2023, **Vol. 20**, p. 1046–1063. DOI: [10.21123/bsj.2023.8227](https://doi.org/10.21123/bsj.2023.8227)
 23. Alsahib S.A., Baqer S.R. Synthesis, identification of some new derivatives of 4-amino antipyrine bearing six and seven membered rings. *Journal of Physics: Conference Series*, 2021, **Vol. 1853**, 012010. DOI: [10.1088/1742-6596/1853/1/012010](https://doi.org/10.1088/1742-6596/1853/1/012010)
 24. Al-Jeilawi O.H., Tuama S.H., Hussein I.A., Shamaya A.N. Synthesis, Characterization, and Biological Evaluation of New Cyclic Quinazoline Derivatives as Potential Antibacterial and Antifungal Agents. *Doklady Chemistry*, 2024, **Vol. 514**, p. 27–34.
 25. Ruaa D.M., Alsahib S.A., Ascar I.F. Synthesis, Characterization and Cyclization of Pyran Using Ag₂O Nanoparticle from Natural Source “Ginger”. *Iraqi Journal of Agricultural Sciences*, 2021, **Vol. 52**, p. 1171–1184. DOI: [10.36103/ijas.v52i5.1455](https://doi.org/10.36103/ijas.v52i5.1455)
 26. Thejeel, K., Abdul Kareem, S., Ascar, I., & Hussein, M. Synthesis of new polymers linked to heterocyclic using zinc oxide with nanostructures extracted from natural sources. *Egyptian Journal of Chemistry*, 2022, **Vol. 65**, p. 579–589. DOI: [10.21608/ejchem.2021.92971.4644](https://doi.org/10.21608/ejchem.2021.92971.4644)
 27. Fahad M.M., Shafiq N., Arshad U., Radh A.J. As Antimicrobial Agents: Synthesis,

- Structural Characterization and Molecular Docking studies of Barbituric Acid Derivatives from Phenobarbital. *Chem. Methodol.* 2022, **Vol. 2**, p. 122-136. DOI:[10.21203/rs.3.rs-1019035/v1](https://doi.org/10.21203/rs.3.rs-1019035/v1)
28. Abd AL-Zahra A.A., Fadel Z.H., Al-Sarray A.J., Hussein I.A., Al-Noor T.H. Corrosion Inhibition of Carbon Steel C45 Using New Azo Derivative in HCl Solution: Synthesis, Potentiostatic Measurement, and DFT Studies. *Russian Journal of Physical Chemistry*, 2024, **Vol. 98**, p. 3202–3211. DOI: [10.1134/S0036024424702674](https://doi.org/10.1134/S0036024424702674)
29. Alsahib S.A., Baqer S.R. Synthesis, identification of some new derivatives of 4-amino antipyrine bearing six and seven membered rings. *Journal of Physics: Conference Series*, 2021, **Vol. 1853**, 012010. DOI: 10.1088/1742-6596/1853/1/012010
30. Ahmed R.S., Ali R.A., Ahamed L.S. Synthesis of new 2,4,5-triphenyl imidazole derivatives derived from benzoin and studying their biological activity. *Journal of Global Pharma Technology*, 2019, **Vol. 11**, p. 437–444.
31. Al-Khazraji S.I., Ahamed L.S., Ali R.A. Synthesis and Characterization of some new heterocyclic derivatives from aromatic carbonyl compounds and carboxylic acids with Evaluation some of them for biological activity. *Iraqi Journal of Science*, 2024, **Vol. 65**, p. 1855–1869. DOI: [10.24996/ij.s.2024.65.4.6](https://doi.org/10.24996/ij.s.2024.65.4.6)
32. Ahamed L.S., Ali R.A., Ahmed R.S. Solvent-free synthesis of new chalcone derivatives from 3-nitro phthalic acid and evaluation of their biological activities. *Egyptian Journal of Chemistry*, 2021, **Vol. 64**, p. 2963–2968. DOI: [10.21608/ejchem.2021.55742.3176](https://doi.org/10.21608/ejchem.2021.55742.3176)
33. Ahamed L.S., Sallal Z.A., Al-Jeilawi O.H., Shamaya A.N., Mahmood A.A.R. Design, Synthesis, and Biological Evaluation of new sulfonamides derived from 2-Aminopyridine: Molecular docking, POM analysis, and identification of the pharmacophore sites. *Moroccan Journal of Chemistry*, 2025, **Vol. 13**, p. 519–539. DOI:[10.48317/IMIST.PRSM/morjchem-v13i2.52998](https://doi.org/10.48317/IMIST.PRSM/morjchem-v13i2.52998)
34. Safi M.N., Ahamed L.S., Almalki F.A., Ben Hadda T. Synthesis, Anticancer, Antimicrobial evaluation, and in Silico Molecular Docking and POM Analyses of New 4,7-dimethyl Coumarin Containing Sulfonamides. *Moroccan Journal of Chemistry*, 2025, **Vol. 13**, p. 849–880. DOI:[10.48317/IMIST.PRSM/morjchem-v13i2.54658](https://doi.org/10.48317/IMIST.PRSM/morjchem-v13i2.54658)
35. Hassan, S. A., Aziz, D. M., Abdullah, M. N., Bhat, A. R., Dongre, R. S., Ben Hadda, T., Almalki, F. A., Kawsar, S. M. A., Rahiman, A. K., Ahmed, S., Abdellattif, M. H., Berredjem, M., Sheikh, S. A., & Jamalis, J. In vitro and in vivo evaluation of the antimicrobial, antioxidant, cytotoxic, hemolytic activities and in silico POM/DFT/DNA-binding and pharmacokinetic analyses of new sulfonamide bearing thiazolidin-4-ones. *Journal of Biomolecular Structure and Dynamics*, **Vol 42**, (2024) 3747–3763. <https://doi.org/10.1080/07391102.2023.2226713>
36. Mohammed A.Y., Ahamed L.S. Synthesis of New Substituted Coumarin Derivatives containing Schiff-Base as Potential Antimicrobial and Antioxidant Agents. *International Journal of Drug Delivery and Technology*, 2022, **Vol. 12**, p. 1279–1281. DOI:10.25258/ijddt.12.3.58